



For women living
with platinum-resistant
ovarian cancer,

Your guide to taking LIFYORLI



What is LIFYORLI?

LIFYORLI is a prescription medicine used with a chemotherapy medicine called nab-paclitaxel to treat adults with ovarian, fallopian tube, or primary peritoneal cancer who:

- Have not responded to or are no longer responding to treatment with platinum-based chemotherapy, and
- Have received 1-3 prior systemic treatments, at least one of which included bevacizumab.

It is not known if LIFYORLI is safe and effective in children.

Please see Important Safety Information on page 11 and Important Facts about LIFYORLI.



Not actual patients



Could LIFYORLI fit into your story?

LIFYORLI plus nab-paclitaxel (a chemotherapy) offers a path forward for women with platinum-resistant ovarian cancer. **You and your healthcare provider can decide together whether this combination is right for you.**

Find important information about:

How to take LIFYORLI

Possible side effects of LIFYORLI plus nab-paclitaxel

Where to find resources and support

Please see [Important Safety Information on page 11](#) and [Important Facts about LIFYORLI](#).

What is LIFYORLI and who is it for?

LIFYORLI is an oral medication taken as three softgel capsules once a day on the day before, the day of, and the day after each nab-paclitaxel infusion. This combination is used to treat adults with ovarian cancer who have not responded to or are no longer responding to treatment with platinum-based chemotherapy, and have received 1-3 prior systemic treatments, at least one of which included bevacizumab.



The packaging shown contains nine doses of LIFYORLI and space for treatment-day reminders on the back.



100-mg
Softgel capsule
Capsules shown are actual size.



25-mg
Softgel capsule



25-mg
Softgel capsule

Take all **three softgel capsules** together with food once a day, preferably in the morning.

CLICK HERE to learn more about how LIFYORLI works in the body

Important Safety Information

Who should not take LIFYORLI?

Do not take LIFYORLI if you are receiving or taking a steroid medicine by mouth or injection (systemic glucocorticoid) for lifesaving purposes. LIFYORLI can make glucocorticoid medicines less effective.


Please see additional Important Safety Information on page 11 and Important Facts about LIFYORLI.

 **Lifyorli™**
(relacorilant) 100 mg, 25 mg capsules

How should I take LIFYORLI?

LIFYORLI is taken as **three softgel capsules once a day** on the **day before**, the **day of**, and the **day after each nab-paclitaxel infusion**. It's important to take LIFYORLI exactly as your healthcare provider tells you.

TREATMENT SCHEDULE



Capsules shown are not actual size.
IV, intravenous.

Important Safety Information (cont'd)

What are the possible side effects of LIFYORLI?

LIFYORLI can cause serious side effects, including:

- **Low white blood cell count (neutropenia) and severe infections.** LIFYORLI in combination with nab-paclitaxel can cause low white blood cell counts, which can lead to a fever and severe infections and cause death. Your healthcare provider will check your blood counts before each weekly treatment with LIFYORLI in combination with nab-paclitaxel and as needed. Tell your healthcare provider right away if you have a fever, chills, weakness, confusion, or other signs or symptoms of an infection.


Please see additional [Important Safety Information on page 11](#) and [Important Facts about LIFYORLI](#).

LIFYORLI plus nab-paclitaxel treatment cycle

The LIFYORLI plus nab-paclitaxel treatment period lasts for **three weeks, followed by one week off**. This is called a cycle, and it repeats every month as needed. You can stay on this treatment schedule as long as you and your doctor feel it's right for you.



During the treatment period, you will receive **a weekly nab-paclitaxel infusion that takes about 30 minutes**.



What is nab-paclitaxel?

Nab-paclitaxel is a type of chemotherapy that is used in combination with LIFYORLI to treat women with platinum-resistant ovarian cancer.


Please see Important Safety Information on page 11 and Important Facts about LIFYORLI.



LIFYORLI comes in blister cards. Each card contains a three-dose supply of LIFYORLI.

CLICK HERE to watch the Taking LIFYORLI video

Helpful tips for taking LIFYORLI



Take it at home with food



Aim to take it at the same time each day, preferably in the morning



Swallow capsules whole. Do not crush, chew, dissolve, or split the capsules



Use your personal planner or set reminders on your phone to track your doses of LIFYORLI

If you vomit after taking LIFYORLI, do not take an extra dose.

What should I do if I miss a dose?

Sometimes things don't go as planned. **If you forget to take LIFYORLI at your regular time, follow these simple guidelines:**



If it has been less than 12 hours

Take it as soon as you remember



If it has been 12 hours or more

Take the next dose at your regular time

Please see [Important Safety Information on page 11](#) and [Important Facts about LIFYORLI](#).

 **Lifyorli™**
(relacorilant) 100 mg, 25 mg capsules

What should I do if the date of my nab-paclitaxel infusion changes?



Stop taking LIFYORLI right away. Start your doses again the day before your next infusion. Take LIFYORLI once the **day before**, the **day of**, and the **day after each nab-paclitaxel infusion**.

Always follow up with your healthcare provider to confirm your new schedule with LIFYORLI. They can also help with any questions or concerns.

Important Safety Information (cont'd)

What are the possible side effects of LIFYORLI? (cont'd)

LIFYORLI can cause serious side effects, including:

- **Reduced effects of the hormone cortisol (adrenal insufficiency).** LIFYORLI reduces the effect of an adrenal hormone called cortisol, which may be life-threatening and require hospitalization. This can happen at any time during treatment. Your risk of adrenal insufficiency is increased when your body is under stress, such as when you are sick, have an infection, or have surgery. Tell your healthcare provider right away if you develop the following symptoms during treatment with LIFYORLI:
 - Feeling dizzy or lightheaded
 - Low blood pressure (hypotension)
 - Low blood sugar (hypoglycemia)
 - Worsening weakness or tiredness
 - Worsening nausea or vomiting
 - Worsening loss of appetite or weight loss

Please see additional Important Safety Information on page 11 and Important Facts about LIFYORLI.

How can I prepare for treatment with LIFYORLI?

It's important to share your complete medical history with your healthcare team. It will help them guide your treatment plan.

Tell your healthcare provider if you take steroid medicines called glucocorticoids



Do not take LIFYORLI if you are receiving or taking a steroid medicine by mouth or injection (systemic glucocorticoid) for lifesaving purposes. LIFYORLI may make steroid medications less effective, and steroids may make LIFYORLI less effective. If you take steroids for another condition, your symptoms may get worse. If you're not sure whether you're taking steroids, ask your healthcare provider.

Share an updated list of your medications



Tell your healthcare provider about all medicines you take, **including prescription and over-the-counter medicines, vitamins, and herbal supplements**. LIFYORLI may affect the way your other medicines work, and other medicines may affect the way LIFYORLI works.

Be prepared for a blood draw



Your healthcare provider will order blood work to **monitor your white blood cell count and other conditions**. Reach out to them if you have any questions about your treatment.

Lead the conversation during your next appointment.

The Care Team Discussion Guide helps ensure your concerns are addressed and your questions are answered.

Please see [Important Safety Information on page 11](#) and [Important Facts about LIFYORLI](#).

 **Lifyorli**[™]
(relacorilant) 100 mg, 25 mg capsules

What are the possible side effects of LIFYORLI plus nab-paclitaxel?

The safety of LIFYORLI was evaluated in a global clinical study of 381 women with platinum-resistant ovarian cancer. Among patients in the study, 9% permanently stopped treatment due to side effects.

Use of LIFYORLI can cause serious side effects including:

Low white blood cell counts (neutropenia) and serious infections, reduced effects of the hormone cortisol (adrenal insufficiency), worsening of conditions treated with glucocorticoids, and harm to an unborn baby.

The most common side effects of LIFYORLI in combination with nab-paclitaxel included:

- Decreased red blood cell, white blood cell, and platelet counts
- Tiredness
- Nausea
- Diarrhea
- Rash
- Decreased appetite

These are not the only possible adverse reactions of LIFYORLI. Call your doctor for medical advice about side effects.

LIFYORLI could harm your unborn baby. It's important that you:

- Take a **pregnancy test** before starting LIFYORLI
- Use **contraception** while on LIFYORLI and for at least one week after your last dose
- **Avoid breastfeeding** during treatment and for at least one week after your last dose



Ask your healthcare provider about effective birth control methods. **Call your healthcare team right away if you become pregnant or think you might be pregnant.**

Please see Important Safety Information on page 11 and Important Facts about LIFYORLI.

 **Lifyorli**[™]
(relacorilant) 100 mg, 25 mg capsules

How do I store LIFYORLI?



Store LIFYORLI at **room temperature** between **68°F to 77°F** (20°C to 25°C)



Be sure to keep LIFYORLI and all medicines **out of the reach of children**

Use your personal planner to manage your treatment plan



Write down your upcoming nab-paclitaxel infusion dates and schedule of LIFYORLI



Keep track of treatment progress and milestones



CLICK HERE
to download extra pages for your planner

Your personal planner will be included in your LIFYORLI Welcome Kit.
If you haven't received yours yet, ask your care team!

Important Safety Information

Who should not take LIFYORLI?

Do not take LIFYORLI if you are receiving or taking a steroid medicine by mouth or injection (systemic glucocorticoid) for lifesaving purposes. LIFYORLI can make glucocorticoid medicines less effective.

What are the possible side effects of LIFYORLI?

LIFYORLI can cause serious side effects, including:

- **Low white blood cell count (neutropenia) and severe infections.**

LIFYORLI in combination with nab-paclitaxel can cause low white blood cell counts, which can lead to a fever and severe infections and cause death. Your healthcare provider will check your blood counts before each weekly treatment with LIFYORLI in combination with nab-paclitaxel and as needed. Tell your healthcare provider right away if you have a fever, chills, weakness, confusion, or other signs or symptoms of an infection.

- **Reduced effects of the hormone cortisol (adrenal insufficiency).**

LIFYORLI reduces the effect of an adrenal hormone called cortisol, which may be life-threatening and require hospitalization. This can happen at any time during treatment. Your risk of adrenal insufficiency is increased when your body is under stress, such as when you are sick, have an infection, or have surgery. Tell your healthcare provider right away if you develop the following symptoms during treatment with LIFYORLI:

- Feeling dizzy or lightheaded
- Worsening weakness or tiredness
- Low blood pressure (hypotension)
- Worsening nausea or vomiting
- Low blood sugar (hypoglycemia)
- Worsening loss of appetite or weight loss

- **Worsening of conditions treated with glucocorticoids.**

Treatment with LIFYORLI in people who also take glucocorticoids for other conditions may worsen symptoms of these conditions. LIFYORLI makes glucocorticoids less effective, and use of glucocorticoids makes LIFYORLI less effective. If you take a glucocorticoid medicine during treatment with LIFYORLI, tell your healthcare provider if symptoms of your other condition get worse.

The most common side effects of LIFYORLI in combination with nab-paclitaxel include

- Decreased red blood cell, white blood cell, and platelet counts
- Tiredness
- Nausea
- Diarrhea
- Rash
- Decreased appetite

Your healthcare provider may change your dose of LIFYORLI, delay treatment, or completely stop treatment with LIFYORLI if you have certain side effects. These are not all the possible side effects of LIFYORLI. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Before you take LIFYORLI, tell your healthcare provider about all your medical conditions, including if you:

- Plan to have surgery or have had surgery within the past 30 days
- Have liver problems
- Are pregnant or plan to become pregnant. LIFYORLI can harm your unborn baby

Females who are able to become pregnant:

- Your healthcare provider should do a pregnancy test before you start treatment with LIFYORLI.
- Use effective birth control (contraception) during treatment with LIFYORLI and for 1 week after your last dose. Talk with your healthcare provider about birth control methods you can use during this time.
- Tell your healthcare provider right away if you become pregnant or think you are pregnant during treatment with LIFYORLI.

Males with female partners who are able to become pregnant:

- Use effective birth control (contraception) during treatment with LIFYORLI and for 1 week after your last dose.
- Are breastfeeding or plan to breastfeed. It is not known if LIFYORLI passes into your breast milk. Do not breastfeed during treatment with LIFYORLI and for 1 week after your last dose.

Tell your healthcare provider about all medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking LIFYORLI with certain other medicines may affect how LIFYORLI or the other medicines work and may cause side effects.

Especially tell your healthcare provider if you are taking steroid medicines called glucocorticoids. Ask your healthcare provider if you are not sure if you take a glucocorticoid.

Please see [Important Facts](#) about LIFYORLI.

 **Lifyorli™**
(relacorilant)
100 mg, 25 mg capsules



Not actual patients

LIFYORLI Support™ is here for you throughout your treatment journey

We can help you:



Understand your insurance coverage



Explore savings options



Get personalized guidance and resources

CLICK HERE

to find support at every step of care

Call LIFYORLI Support at **1-85-LIFYORLI (1-855-439-6754)**,
Monday through Friday, 8 AM to 8 PM ET,
or visit LIFYORLI.com/support.

Please see Important Safety Information on page 11 and Important Facts about LIFYORLI.

 **Lifyorli™**
(relacorilant) 100 mg, 25 mg
capsules

 **Corcept**
THERAPEUTICS

LIFYORLI, LIFYORLI Support, and the LIFYORLI logo are trademarks of Corcept. Corcept and the Corcept logo are registered trademarks. © 2026 Corcept. US-REL-26-0102 04-26