

IMPORTANT FACTS

This is only a brief summary of important information about LIFYORLI. Talk to your healthcare provider or pharmacist to learn more.

About LIFYORLI (lif-YOR-lee)

LIFYORLI is a prescription medicine used with a chemotherapy medicine called nab-paclitaxel to treat adults with ovarian, fallopian tube, or primary peritoneal cancer who:

- Have not responded to or are no longer responding to treatment with platinum-based chemotherapy, and
- Have received 1-3 prior systemic treatments, at least one of which included bevacizumab

It is not known if LIFYORLI is safe and effective in children.

Do not take LIFYORLI if you are receiving or taking a steroid medicine by mouth or injection for lifesaving purposes.

Before taking LIFYORLI (cont'd)

Males with female partners who are able to become pregnant:

- Use effective birth control (contraception) during treatment with LIFYORLI and for 1 week after your last dose
- Are breastfeeding or plan to breastfeed. It is not known if LIFYORLI passes into your breast milk. Do not breastfeed during treatment with LIFYORLI and for 1 week after your last dose

Tell your healthcare provider about all

medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking LIFYORLI with certain other medicines may affect how LIFYORLI or the other medicines work and may cause side effects.

Especially tell your healthcare provider if you are taking steroid medicines called glucocorticoids.

Ask your healthcare provider if you are not sure if you take a glucocorticoid.

Before taking LIFYORLI

Before you take LIFYORLI, tell your healthcare provider about all your medical conditions, including if you:

- Plan to have surgery or have had surgery within the past 30 days
- Have liver problems
- Are pregnant or plan to become pregnant. LIFYORLI can harm your unborn baby

Females who are able to become pregnant:

- Your healthcare provider should do a pregnancy test before you start treatment with LIFYORLI
- Use effective birth control (contraception) during treatment with LIFYORLI and for 1 week after your last dose. Talk with your healthcare provider about birth control methods you can use during this time
- Tell your healthcare provider right away if you become pregnant or think you are pregnant during treatment with LIFYORLI

Possible side effects of LIFYORLI + nab-paclitaxel

LIFYORLI can cause serious side effects, including:

- **Low white blood cell count (neutropenia) and severe infections.** LIFYORLI in combination with nab-paclitaxel can cause low white blood cell counts, which can lead to a fever and severe infections and cause death. Your healthcare provider will check your blood counts before each weekly treatment with LIFYORLI in combination with nab-paclitaxel and as needed. Tell your healthcare provider right away if you have a fever, chills, weakness, confusion, or other signs or symptoms of an infection

IMPORTANT FACTS (cont'd)

Possible side effects of LIFYORLI + nab-paclitaxel (cont'd)

- **Reduced effects of the hormone cortisol (adrenal insufficiency).** LIFYORLI reduces the effect of an adrenal hormone called cortisol, which may be life-threatening and require hospitalization. This can happen at any time during treatment. Your risk of adrenal insufficiency is increased when your body is under stress, such as when you are sick, have an infection, or have surgery. Tell your healthcare provider right away if you develop the following symptoms during treatment with LIFYORLI:
 - Feeling dizzy or lightheaded
 - Low blood pressure (hypotension)
 - Low blood sugar (hypoglycemia)
 - Worsening weakness or tiredness
 - Worsening nausea or vomiting
 - Worsening loss of appetite or weight loss
- **Worsening of conditions treated with glucocorticoids.** Treatment with LIFYORLI in people who also take glucocorticoids for other conditions may worsen symptoms of these conditions. LIFYORLI makes glucocorticoids less effective, and use of glucocorticoids makes LIFYORLI less effective. If you take a glucocorticoid medicine during treatment with LIFYORLI, tell your healthcare provider if symptoms of your other condition get worse

The most common side effects of LIFYORLI in combination with nab-paclitaxel include:

- Decreased red blood cell, white blood cell, and platelet counts
- Tiredness
- Nausea
- Diarrhea
- Rash
- Decreased appetite

Your healthcare provider may change your dose of LIFYORLI, delay treatment, or completely stop treatment with LIFYORLI if you have certain side effects.

These are not the only possible side effects of LIFYORLI. Call your healthcare provider for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How to take LIFYORLI

- Take LIFYORLI exactly as your healthcare provider tells you
- Follow the LIFYORLI dosing instructions on the blister card
- Take LIFYORLI 1 time on the **day before**, the **day of**, and the **day after** each nab-paclitaxel infusion
- Take LIFYORLI with food
- Swallow LIFYORLI capsules whole. Do not crush, chew, dissolve, or split the capsules
- If you miss a dose of LIFYORLI
 - **By less than 12 hours**, take it as soon as you remember
 - **By more than 12 hours**, skip the missed dose and take the next dose at your regular time
 - **Do not** take 2 doses at the same time to make up for a missed dose
- If you vomit after taking LIFYORLI, **do not** take an extra dose

Get more information

- This is only a brief summary of important information about LIFYORLI. Talk to your healthcare provider or pharmacist to learn more
- To access LIFYORLI Support™, visit LIFYORLI.com/support or call **1-85-LIFYORLI (1-855-439-6754)**

[CLICK HERE](#) to see additional Important Safety Information

