



Frequently Asked Questions

Find answers to common questions about taking LIFYORLI plus nab-paclitaxel for platinum-resistant ovarian cancer.



Not an actual patient

What is LIFYORLI?

LIFYORLI is a prescription medicine used with a chemotherapy medicine called nab-paclitaxel to treat adults with ovarian, fallopian tube, or primary peritoneal cancer who:

- Have not responded to or are no longer responding to treatment with platinum-based chemotherapy, and
- Have received 1-3 prior systemic treatments, at least one of which included bevacizumab.

It is not known if LIFYORLI is safe and effective in children.

Important Safety Information

Who should not take LIFYORLI?

Do not take LIFYORLI if you are receiving or taking a steroid medicine by mouth or injection (systemic glucocorticoid) for lifesaving purposes. LIFYORLI can make glucocorticoid medicines less effective.

Please see additional Important Safety Information and Important Facts about LIFYORLI.

About LIFYORLI

What is LIFYORLI and who is it for?

LIFYORLI is a medication taken as three softgel capsules along with nab-paclitaxel (a chemotherapy). It is used to treat adults with ovarian cancer who:

- Have not responded to or are no longer responding to treatment with platinum-based chemotherapy, and
- Have received 1-3 prior systemic treatments, at least one of which included bevacizumab.

It is not known if LIFYORLI is safe and effective in children.

Do not take LIFYORLI if you are receiving or taking a steroid medicine by mouth or injection (systemic glucocorticoid) for lifesaving purposes. LIFYORLI can make glucocorticoid medicines less effective.

Do I need a genetic test or other type of test to take LIFYORLI?

No, you don't need to take any specific test (such as a genetic or biomarker test—which helps identify a gene or protein in your cancer cells) to be treated with LIFYORLI.

What results could I expect with LIFYORLI?

LIFYORLI was evaluated in a global clinical study of 381 patients called ROSELLA. Compared to nab-paclitaxel alone, LIFYORLI plus nab-paclitaxel helped women live longer and reduced the risk of cancer getting worse.* Specifically, the study showed:

- **35% reduction in the risk of death**
- **30% lower risk of cancer progression***

Please remember that everyone's experience is different, so results may vary.

*Progression was defined as the cancer getting worse, as measured by scans, or if the patient died for any reason, whichever occurred first.

Learn more about how LIFYORLI was studied

SEE RESULTS WITH LIFYORLI

Please see [Important Safety Information](#) and [Important Facts](#) about LIFYORLI.

Taking LIFYORLI + nab-paclitaxel

How should I take LIFYORLI?

LIFYORLI is taken as **three softgel capsules once a day** on the **day before**, the **day of**, and the **day after each nab-paclitaxel infusion**. Aim to take it at the same time each day, preferably in the morning.

Take LIFYORLI with food. You should swallow the capsules whole, and do not crush, chew, dissolve, or split the capsules. If you vomit after taking LIFYORLI, **do not** take an extra dose.

The LIFYORLI plus nab-paclitaxel cycle is given for three weeks, followed by one week off.

It's important to take LIFYORLI exactly as your healthcare provider tells you.

How is nab-paclitaxel administered?

Nab-paclitaxel is a treatment administered by a healthcare provider. **It is given by infusion, which typically lasts 30 minutes.** Nab-paclitaxel is administered once a week for three weeks in a row, followed by a rest week.

Are there medications I should avoid while taking LIFYORLI?

Yes. You should tell your healthcare provider about all medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. LIFYORLI may affect the way your other medicines work, and other medicines may affect the way LIFYORLI works.

Especially tell your healthcare provider if you are taking steroid medicines called glucocorticoids. If you're not sure whether you're taking steroids, ask them.

What happens if I miss a dose of LIFYORLI?

If you forget to take LIFYORLI at your regular time, follow these simple guidelines:

- **If it has been less than 12 hours**, take it as soon as you remember
- **If it has been 12 hours or more**, take the next dose at your regular time

Contact your healthcare provider with any questions or concerns.

How should I adjust my LIFYORLI doses if the date of my nab-paclitaxel infusion changes?

Stop taking LIFYORLI right away and restart your once-daily doses the day before your new scheduled infusion. Take your medicine the day before, the day of, and the day after each nab-paclitaxel infusion.

Follow up with your healthcare provider to confirm your new schedule with LIFYORLI.

Side effects with LIFYORLI + nab-paclitaxel


What are the possible side effects of LIFYORLI plus nab-paclitaxel?

Use of LIFYORLI can cause serious side effects including: low white blood cell counts (neutropenia) and serious infections, reduced effects of the hormone cortisol (adrenal insufficiency), worsening of conditions treated with glucocorticoids, and harm to an unborn baby.

The most common side effects of LIFYORLI plus nab-paclitaxel are consistent with those associated with nab-paclitaxel alone, including:

- Decreased red blood cell, white blood cell, and platelet counts
- Tiredness
- Nausea
- Diarrhea
- Rash
- Decreased appetite

These are not the only possible side effects of LIFYORLI. Follow up with your healthcare provider about any side effects.



Talk to your healthcare provider to see if LIFYORLI might be right for you

You can also download a helpful brochure to learn more about LIFYORLI.

GET THE BROCHURE

Not an actual patient

Please see [Important Safety Information](#) and [Important Facts](#) about LIFYORLI.

What kind of support services are available when taking LIFYORLI?

LIFYORLI Support can help you get started by connecting you with:

- **Case Managers** who can help you understand your insurance coverage and determine your potential out-of-pocket costs for LIFYORLI
- **LIFYORLI Support Nurse Educators** to answer questions you may have about LIFYORLI*

To learn more, call us at 1-85-LIFYORLI **(1-855-439-6754)**, Monday through Friday, 8 AM to 8 PM ET, or visit LIFYORLI.com/support.

Are there financial programs available to help me afford LIFYORLI?

- If you have commercial (also known as private) health insurance, you may be eligible to **pay as little as \$0** per month for LIFYORLI†
- LIFYORLI Support can help you identify options for financial assistance, depending on your insurance type

To learn more, call us at 1-85-LIFYORLI **(1-855-439-6754)**, Monday through Friday, 8 AM to 8 PM ET, or visit LIFYORLI.com/support.

I'm a caregiver for someone who was prescribed LIFYORLI. Can I also receive information and resources from LIFYORLI Support?

Yes! You're a part of your loved one's support team throughout their treatment journey, and **we want you to feel supported, too.**

Explore LIFYORLI.com/support to learn what LIFYORLI Support offers you and your loved one. You can also call us at 1-85-LIFYORLI **(1-855-439-6754)**, Monday through Friday, 8 AM to 8 PM ET.

*A LIFYORLI Support Nurse Educator cannot provide medical or clinical advice; these questions should be directed to your doctor, nurse, or pharmacist.

†For full Terms and Conditions, please visit LIFYORLIsavingscard.com.

Important Safety Information

Who should not take LIFYORLI?

Do not take LIFYORLI if you are receiving or taking a steroid medicine by mouth or injection (systemic glucocorticoid) for lifesaving purposes. LIFYORLI can make glucocorticoid medicines less effective.

What are the possible side effects of LIFYORLI?

LIFYORLI can cause serious side effects, including:

- **Low white blood cell count (neutropenia) and severe infections.** LIFYORLI in combination with nab-paclitaxel can cause low white blood cell counts, which can lead to a fever and severe infections and cause death. Your healthcare provider will check your blood counts before each weekly treatment with LIFYORLI in combination with nab-paclitaxel and as needed. Tell your healthcare provider right away if you have a fever, chills, weakness, confusion, or other signs or symptoms of an infection.
- **Reduced effects of the hormone cortisol (adrenal insufficiency).** LIFYORLI reduces the effect of an adrenal hormone called cortisol, which may be life-threatening and require hospitalization. This can happen at any time during treatment. Your risk of adrenal insufficiency is increased when your body is under stress, such as when you are sick, have an infection, or have surgery. Tell your healthcare provider right away if you develop the following symptoms during treatment with LIFYORLI:
 - Feeling dizzy or lightheaded
 - Low blood pressure (hypotension)
 - Low blood sugar (hypoglycemia)
 - Worsening weakness or tiredness
 - Worsening nausea or vomiting
 - Worsening loss of appetite or weight loss
- **Worsening of conditions treated with glucocorticoids.** Treatment with LIFYORLI in people who also take glucocorticoids for other conditions may worsen symptoms of these conditions. LIFYORLI makes glucocorticoids less effective, and use of glucocorticoids makes LIFYORLI less effective. If you take a glucocorticoid medicine during treatment with LIFYORLI, tell your healthcare provider if symptoms of your other condition get worse.

The most common side effects of LIFYORLI in combination with nab-paclitaxel include

- Decreased red blood cell, white blood cell, and platelet counts
- Tiredness
- Nausea
- Diarrhea
- Rash
- Decreased appetite



Your healthcare provider may change your dose of LIFYORLI, delay treatment, or completely stop treatment with LIFYORLI if you have certain side effects. These are not all the possible side effects of LIFYORLI. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Before you take LIFYORLI, tell your healthcare provider about all your medical conditions, including if you:

- Plan to have surgery or have had surgery within the past 30 days
- Have liver problems
- Are pregnant or plan to become pregnant. LIFYORLI can harm your unborn baby

Females who are able to become pregnant:

- Your healthcare provider should do a pregnancy test before you start treatment with LIFYORLI.
- Use effective birth control (contraception) during treatment with LIFYORLI and for 1 week after your last dose. Talk with your healthcare provider about birth control methods you can use during this time.
- Tell your healthcare provider right away if you become pregnant or think you are pregnant during treatment with LIFYORLI.

Males with female partners who are able to become pregnant:

- Use effective birth control (contraception) during treatment with LIFYORLI and for 1 week after your last dose.
- Are breastfeeding or plan to breastfeed. It is not known if LIFYORLI passes into your breast milk. Do not breastfeed during treatment with LIFYORLI and for 1 week after your last dose.

Tell your healthcare provider about all medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking LIFYORLI with certain other medicines may affect how LIFYORLI or the other medicines work and may cause side effects.

Especially tell your healthcare provider if you are taking steroid medicines called glucocorticoids. Ask your healthcare provider if you are not sure if you take a glucocorticoid.

Please see [Important Facts](#) about LIFYORLI.

LIFYORLI, LIFYORLI Support, and the LIFYORLI logo are trademarks of Corcept. Corcept and the Corcept logo are registered trademarks.
© 2026 Corcept. US-REL-25-0029 03-26

